

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004-2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG):

R9BKBE

Southeast Gulf Coastal Plain Blackland Prairie and Woodland

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Will McDearman

will_mcdearman@fws.gov

Malcolm Hodges

mhodges@tnc.org

Reviewers

Vegetation Type

Grassland

Dominant Species*

| | |
|-------|--------|
| SCSC | ANGE |
| SONU2 | QUMA3 |
| JUVI | TRDA3 |
| QUST | SPCO16 |

General Model Sources

- Literature
- Local Data
- Expert Estimate

LANDFIRE Mapping Zones

| |
|----|
| 46 |
| 55 |

Rapid Assessment Model Zones

- | | |
|--|---|
| <input type="checkbox"/> California | <input type="checkbox"/> Pacific Northwest |
| <input type="checkbox"/> Great Basin | <input type="checkbox"/> South Central |
| <input type="checkbox"/> Great Lakes | <input checked="" type="checkbox"/> Southeast |
| <input type="checkbox"/> Northeast | <input type="checkbox"/> S. Appalachians |
| <input type="checkbox"/> Northern Plains | <input type="checkbox"/> Southwest |
| <input type="checkbox"/> N-Cent. Rockies | |

Geographic Range

Black Belt or Blackland Prairie occurs mainly in the gulf coastal plain of Tennessee, Mississippi and Alabama in a broad arc approximately 450 km long by 40-50 km wide (Subsection 231Ba of Keys et al. 1995; Ecoregion 65a of Griffith et al. 2001). The system also includes the Jackson Prairie region of Mississippi; the Chunnenugee Hills, Red Hills, and Lime Hills of Alabama in Washington, Wilcox, Monroe and Clark counties; and prairie remnants in the Fort Valley Plateau of Bleckley and Houston counties, Georgia (FRCC model, NatureServe 2005).

Biophysical Site Description

Blackland prairie and woodland occurs on eponymous rich, black, circumneutral topsoils formed over clayey, heavy, usually calcareous subsoils with carbonatic or montmorillonitic mineralogy. The system occurs in association with formations of the Tertiary Jackson (Yazoo Clay), Claiborne (Cook Mountain) and Fleming groups; and the Cretaceous Selma group (Selma, Mooreville or Demopolis chalks). The matrix around the blackland prairies is pine-oak forests growing in acidic, sandier soils with less clay (recent STATSCO soils maps).

Floristic similarity among sites across this geographic range generally appears to be 50% or greater, although a number of different alliances within this type have been recognized according to dominant, co-dominant, and diagnostic species. Extant prairies occur in single patches as well as mosaics less than one acre to over several hundred acres in response to soil depth, slope and fire. Mosaics may include virtually treeless patches associated with other patches of widely scattered trees, open deciduous woodlands and evergreen thickets (red cedar "balds").

This PNVG represents a mosaic of Eastern Red cedar Woodland, and Post Oak – Blackjack Oak Woodland, and Little Bluestem – Yellow Indiagrass Herbaceous alliances, as classified by NatureServe

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

(2005). It is a rare and imperiled vegetation type consisting of scattered remnants. Most of the original cover has been destroyed or altered by conversion to agriculture and the exclusion of fire.

Vegetation Description

Blackland prairie and woodland is a mosaic of southeastern dry-mesic tallgrass vegetation, deciduous woodlands and red cedar thickets or "balds." The prairies are united by the relative abundance of little bluestem (*Schizachyrium scoparium*) and yellow Indian grass (*Sorghastrum nutans*), with other herbs including big bluestem (*Andropogon gerardii*), eastern gamma grass (*Tripsacum dactyloides*), composite dropseed (*Sporobolus compositus*), sideoats grama (*Bouteloua curtipendula*), white prairie clover (*Dalea candida*), purple prairie clover (*Dalea purpurea*), pale purple coneflower (*Echinacea pallida*), blazing-star (*Liatris* spp.), and rosin-weed (*Silphium*). Woody species include post oak (*Quercus stellata*), blackjack oak (*Q. marilandica*), chinquapin oak (*Q. muhlenbergii*), black oak (*Q. velutina*), southern red oak (*Q. falcata*), Durand oak (*Q. sinuata* var. *sinuata*), American elm (*Ulmus americana*), green ash (*Fraxinus pennsylvanica*), smooth sumac (*Rhus glabra*), winged sumac (*Rhus copallina*), and eastern red cedar (*Juniperus virginiana*).

Disturbance Description

For the last 500-1000 years, fires were probably annual in most of the system, many if not most set by aboriginals. Fires were probably used to clear prairies for agricultural planting, to eliminate woody growth, and to aid in hunting. The modern landscape shows a tendency toward erosion, creating shallow-soil areas known as "cedar balds" where soil erosion, presumably from historic agriculture or over-grazing, has reduced topsoil. These areas often show exposures of underlying chalk. Such areas may have resulted (albeit at much lower frequencies) from aboriginal agriculture or overgrazing by native herbivores.

Adjacency or Identification Concerns

Vegetation differences from mosaic pine-oak forests are dramatic, and can be discerned at a glance from the trained eye even in heavily disturbed sites by looking for key dominant woody plant species. Soils maps show higher clay content and lower sand percentages than for surrounding areas (recent STATSCO soils maps).

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Disturbance occurred at a scale often larger than the size of patches, although a significant portion probably originated in the present system. These prairie-woodland mosaics occurred in a pyric matrix, so fire probably swept in frequently from outside. Patches on the order of 40-50 square miles may have burned following a well planned ignition, assuming nocturnal humidity recovery completely extinguished the fire. Smaller patches would have burned within natural barriers, different ignition parameters, and less extreme weather.

Issues/Problems

This model is inclusive, with related systems such as Jackson Prairie and Georgia outliers added to the typical Black Belt type. This makes description of the system less definitive and possibly confusing. The characterization of the type as a woodland - prairie mosaic rather than a simple prairie (as was done in the original PNVG) is truer to what probably occurred pre-historically, but makes the system more difficult to model and describe.

Model Evolution and Comments

PNVG Code: BKBE. Hodges borrowed heavily from McDearman's PNVG description, but changed McDearman's VDDT model to include woodland mosaic from the beginning, and to include an optional disturbance, erosion of topsoil to form red cedar balds (Optional 1 disturbance).

One anonymous review was completed for this model. The reviewer stated the model and description

should be accepted as is.

Succession Classes

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 70%

Early1 Open

Description

Fire is almost annual in Class A. Large patches of prairie grasslands are the dominant ecotype, covering an estimated 50-90% of the landscape. Mosaics of prairie, thicket and woodland are dominated by prairies in uplands, in a complex, dendritic landscape, with heterogeneous age classes in woodlands and interconnected grassland patches. Red cedar at the edges of bald thickets are scorched or killed by fire. With no fire, Class A disappears within 5 years in this model, when it succeeds to Class B.

Indicator Species* and Canopy Position

SCSC Lower
SONU2 Lower
ANGE Lower
TRDA3 Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 3

Structure Data (for upper layer lifeform)

| | Min | Max |
|-----------------|------------------|----------------|
| Cover | 0 % | 60 % |
| Height | Herb Short <0.5m | Herb Tall > 1m |
| Tree Size Class | no data | |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 25%

Mid1 Open

Description

Fire has been excluded for 5 years. Large patches of prairie are uncommon to rare, covering about 25-50% of the landscape. Mosaics tend toward dominance by deciduous woodlands, with some prairie patches interconnected and some isolated within woodlands. Forests are rare. With no fire, Class B disappears within 25 years in this model, succeeding to Class C. With fire, Class B still has enough fuels to cycle back to Class A; I postulate no closed loop perpetuating Class B.

Indicator Species* and Canopy Position

SCSC Lower
JUVI Middle
QUST Mid-Upper
SONU2 Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 2

Structure Data (for upper layer lifeform)

| | Min | Max |
|-----------------|-----------------|--------------------|
| Cover | 0 % | 60 % |
| Height | no data | Tree Medium 10-24m |
| Tree Size Class | Medium 9-21"DBH | |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Grasses remain growing under the tree cover in this transitional class.

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 0%

Mid1 All Structures

Description

Fire has been excluded for 25 years. Woodlands are the dominant cover type, with prairie patches generally isolated or small, covering about 5-25% of the landscape. The woodland canopy cover is taller, with trees tending toward maturity. Some patches of forest are present. Species within forests tend toward a mix of fire tolerant and intolerant. With no fire, Class C disappears within 50 years in this model, succeeding to Class D. Class C might have persisted in somewhat isolated situations with occasional fire (25-50-year return interval).

Indicator Species* and Canopy Position

QUST Upper
JUVI Middle
QUMA3 Middle
SCSC Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 2

Structure Data (for upper layer lifeform)

| | Min | Max |
|-----------------|-----------------|--------------------|
| Cover | 30 % | 100 % |
| Height | no data | Tree Medium 10-24m |
| Tree Size Class | Medium 9-21"DBH | |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 0%

Late1 Closed

Description

Fire has been excluded for 50+ years. Forests are dominant, with patches of woodland and rare prairie remnants surviving only in edaphically controlled conditions in less than 5% of the landscape. Forests are mixed, with fire-intolerant species achieving dominance in many areas. With no fire, Class D dominates the landscape, with 95% coverage after 50 years, according to this model. This could happen in a forest patch isolated by wetlands.

Indicator Species* and Canopy Position

QUST Upper
CELA Upper
FRVI Upper
JUVI Mid-Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 8

Structure Data (for upper layer lifeform)

| | Min | Max |
|-----------------|-----------------|------------------|
| Cover | 60 % | 100 % |
| Height | no data | Tree Tall 25-49m |
| Tree Size Class | Large 21-33"DBH | |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class E 5%

Late2 All Structures

Description

Erosion has removed most or all topsoil from subsoils or Cretaceous chalk. Eastern red cedar occurs in patches of varying size, from individuals through small groups to large thickets. The canopy is variable, with patches of trees interspersed with bare ground or sparse herb cover. Outcroppings of chalk occur. Fire may remove some or all red cedars in a patch, but the class depends on lack of soil, so the presumption is that it will reseed and replace itself. This Class may have resulted from aboriginal agriculture or via overgrazing by native grazers. I assume that if left alone, soils will form as plants colonize the patch, and that Class E will eventually succeed to Class A. A default of 500 years is used for this transition, but this is a guess.

Indicator Species* and Canopy Position

JUVI Low-Mid
RHGL Low-Mid
RHCO Lower
SCSC Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 4

Structure Data (for upper layer lifeform)

| | Min | Max |
|-----------------|-----------------|--------------------|
| Cover | 25 % | 85 % |
| Height | no data | Tree Medium 10-24m |
| Tree Size Class | Medium 9-21"DBH | |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

Non-Fire Disturbances Modeled

- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other: Aboriginal agric. & grazing
- Other:

Fire Regime Group: 1

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Historical Fire Size (acres)

Avg:
Min:
Max:

Fire Intervals (FI):

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

| | Avg FI | Min FI | Max FI | Probability | Percent of All Fires |
|-------------|--------|--------|--------|-------------|----------------------|
| Replacement | 7 | | | 0.14286 | 24 |
| Mixed | 2.2 | | | 0.45455 | 76 |
| Surface | | | | | |
| All Fires | 2 | | | 0.59741 | |

*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

References

- Brown, A.J. 1894. History of Newton County from 1834-1894. Republished by Melvin Tingle, Decatur, MS. Fulton, MS: Itawamba County Times, Inc.
- Brown, James K. and Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT.: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.
- DeSelm, H.R. and Murdock, N. 1993. Grass-dominated communities. In. Martin, W.H., Boyce, S.G. and Echternacht, A.C., ed. Biodiversity of the southeastern United States. New York, NY: John Wiley and Sons. Pp 87-141.
- Elsen, D. and Wieland, R. 2003. Rediscovery and management of prairie remnants of the Bienville National Forest, east-central Mississippi. In Peacock E. and Schauwecker, T., ed. Blackland prairies of the Gulf coastal plain. Tuscaloosa, AL: University of Alabama Press, pp. 240-245.
- Feathersonhaugh, G.W. 1844. Geological report of an examination made in 1834 of the elevated country between the Missouri and Red Rivers. Ex-Doc. 152, 23rd Cong. 2nd session. Vol. IV. 97pp.
- Foti, T.L. 1989. Blackland prairies of southwestern Arkansas. Proceedings of the Arkansas Academy of Science 43:23-28.
- Foti, T.L., Simon, S., Zollner, D. and Hattengach, M. 2003. Blackland prairie landscapes of southwestern Arkansas. In Peacock, E. and Schauwecker, T. ed. Blackland prairies of the Gulf coastal plain. Tuscaloosa, AL: University of Alabama Press. Pp. 95-109.
- Gibson, D.J. and Hubert, L.C. 1987. Effects of fire, topography and year-to-year climatic variation on species composition in tallgrass prairie. *Vegetation* 72:175-185.
- Gordon, K.L. and Wiseman, J.B., Jr. 1989. Bienville National Forest prairie survey. Museum Technical Report No. 7. Jackson, MS: Mississippi Department of Wildlife, Fisheries and Parks – Mississippi Museum of Natural Science.
- Harper, R.M. 1913. A botanical cross-section of northern Mississippi. *Bulletin of the Torrey Botanical Club* 40:377-399.
- Harper, R.M. 1920. The limestone prairies of Wilcox County, Alabama. *Ecology* 1:198-213.
- Hilgard, E. 1860. Report on the geology and agriculture of the state of Mississippi. Jackson, MS: E. Banksdale. State Printer. 391 pp.
- Jones, A.S. and Patton, E.G. 1966. Forest, "prairie" and soils in the Black Belt of Sumter County, Alabama in 1832. *Ecology* 47:75-80.
- Jones, S.A. 1971. A virgin prairie and a virgin loblolly pine stand in central Mississippi. *Castanea* 36:223-225.
- Leidolf, A. and McDaniel, S. 1998. A floristic study of black prairie plant communities at Sixteen Section Prairie, Oktibbeha County, Mississippi. *Castanea* 63:51-62.

Lowe, E.N. 1913. Notes on the flora of Mississippi. Mississippi Geological Survey Bulletin No. 12. Jackson, MS: Bureau of Geology. 335 pp.

MacRoberts, M.H., MacRoberts, B.R. and Jackson, L.S. 2003. Louisiana prairies. In Peacock, E. and Schauwecker, T., ed. Blackland prairies of the Gulf coastal plain. Tuscaloosa, AL: University of Alabama Press, pp 81-93.

Moran, L.P. 1995. Soil and ecology of prairie remnants in the Jackson Prairie Region of Mississippi. M.S. Thesis. Mississippi State, MS: Mississippi State University.

Moran, L.P., Pettry, D.E., Switzer, R.E., McDaniel, S.T. and Wieland, R. 1997. Soils of native prairie remnants in the Jackson Prairie region of Mississippi. Bulletin 1067. Mississippi State, MS: Mississippi Agriculture and Forestry Experiment Station.

Myers, M.W. 1948. Geography of the Mississippi Black Prairie. Ph.D. dissertation. Worcester, MA: Clark University.

Nuzzo, V.A. 1986. Extent and status of Midwest oak savanna: presettlement and 1985. *Natural Areas Journal* 6:6-36.

Rankin, H.T. and D.E. Davis. 1971. Woody vegetation in the Black Belt prairie of Montgomery County, Alabama in 1845-46. *Ecology* 52:716-719.

Rostlund, E. 1957. The myth of a natural prairie belt in Alabama: an interpretation of historical records. *Annals of the American Association of Geographers* 47:392-411.

Schauwecker, T. and MacDonald, J. 2003. Plant assemblage response to disturbance at a blackland prairie restoration site in northeastern Mississippi. In Peacock, E. and Schauwecker, T., ed. Blackland prairies of the Gulf coastal plain. Tuscaloosa, AL: University of Alabama Press. Pp. 247-253.

Schmidt, Kirsten M., Menakis, James P., Hardy, Colin C., Hann, Wendel J., and Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

Schuster, M.F. and McDaniel, S. 1973. A vegetation analysis of a black prairie relict site near Aliceville, Alabama. *Journal of the Mississippi Academy of Science* 19:153-159.

U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/>.

Wackerman, A.E. 1929. Why prairies in Arkansas and Louisiana? *Journal of Forestry* 27:726-734.

Wailles, L.C. 1854. Report on the agriculture and geology of Mississippi: embracing a sketch of the social and natural history of the state. Jackson, MS: Mississippi Legislature.

Weiher, E., Forbes, S., Schauwecker, S. and Grace, B. 2004. Multivariate control of plant species richness

and community biomass in blackland prairie. *Oikos* 106:151-157.

Wieland, R.G. 1994. Preliminary vegetation survey and analysis of the Chickasaw Village Site, Natchez Trace Parkway, Tupelo, Mississippi. Museum Technical Report No. 26. Jackson, MS: Mississippi Department of Wildlife, Fisheries and Parks, Mississippi Museum of Natural Science.

Wieland, R.G. 1994. Mississippi Natural Heritage Program: Ecological Communities. Unpublished. Jackson, MS: Mississippi Department of Wildlife, Fisheries and Parks, Museum of Natural Science, Natural Heritage Program. 7 p.

Wieland, R.G. 1995. Jackson Prairie openings, clay barrens of the Gulf Coastal Plain. Jackson, MS: Mississippi Department of Wildlife, Fisheries and Parks, Museum of Natural Science, Natural Heritage Program.

Wieland, R.G., Gordon, K.L., Wiseman, J.B. and Elsen, D.S. 1991. Agencies inventory and restore prairie openings in Bienville National Forest (Mississippi). *Restoration and Management Notes* 9:105-106.

Zollner, D., Simon, S. and Foti, T. L. 2003. A plant community classification for Arkansas's Blackland Prairie Ecosystem. In Peacock, E. and Schauwecker, T., eds. *Blackland prairies of the Gulf coastal plain*. Tuscaloosa, AL: University of Alabama Press, pp. 111-145.

PERSONAL COMMUNICATION (if applicable):

Tom Foti, Arkansas Natural Heritage Commission, Little Rock, AR.

Ron Masters, Tall Timbers Research Station, Tallahassee, FL.

Ron Wieland, Mississippi Department of Wildlife, Fisheries and Parks, MS Museum of Natural Science, Jackson, MS.

Douglas Zollner, The Nature Conservancy, Arkansas Chapter, Little Rock, AR.

Latimore Smith, The Nature Conservancy, Louisiana Chapter, Baton Rouge, LA.

Hodges' references:

Griffith, G.E., Omernik, J.M., Comstock, J.A., Lawrence, S., Martin, G., Goddard, A., Hulcher, V.J. and Foster, T. 2001. *Ecoregions of Alabama and Georgia*. (Two-sided color poster with map, descriptive text, summary tables, and photographs). Reston, VA: U.S. Geological Survey. Scale 1:1,700,000.

Keys, J.E., Jr., Carpenter, C.A., Hooks, S.L., Koenig, F.G., McNab, W.H., Russell, W.E. and Smith, M.L. 1995. *Ecological units of the eastern United States - first approximation* (map and booklet of map unit tables). Presentation scale 1:3,500,000, colored. Atlanta, GA: USDA Forest Service.

NatureServe. 2005. *International Ecological Classification Standard: Terrestrial Ecological Classifications*. Terrestrial ecological systems of the Southeast Region US: DRAFT legend for Landfire project. Arlington, VA: NatureServe Central Databases. Data current as of 25 February 2005.

STATSCO maps courtesy of Jim Menakis, LandFire workshop cadre.